



Pediatric Fever Management

You may alternate Tylenol and Ibuprofen every 3-4 hours as needed for fever in children older than 6 months. Ibuprofen should not be given to patients under 6 months of age.

Children's TYLENOL Dosage Schedule

Weight	Infant Drops 80 mg/0.8 mL	Suspension 160 mg/5 mL
6-11 lbs	½ dropper 0.4 mL	
12-17 lbs	1 dropper 0.8 mL	½ teaspoon 80 mg
18-23 lbs	1 ½ dropper 1.2 mL	¾ teaspoon 120 mg
24-35 lbs	2 droppers 1.6 mL	1 teaspoon 160 mg
36-47 lbs		1 ½ teaspoon 240 mg
48-59 lbs		2 teaspoons 320 mg
60-71 lbs		2 ½ teaspoons 400 mg
72-95 lbs		3 teaspoons 480 mg
96 - > lbs		4 teaspoons 640 mg

Children's IBUPROFEN Dosage Schedule

Suspension 100 mg/5 mL

Weight	<102.5 F	>102.5 F
13-17 lbs	¼ teaspoon 25 mg	½ teaspoon 50 mg
18-23 lbs	½ teaspoon 50 mg	1 teaspoon 100 mg
24-35 lbs	¾ teaspoon 75 mg	1 ½ teaspoon 150 mg
36-47 lbs	1 teaspoon 100 mg	2 teaspoon 200 mg
48-59 lbs	1 ¼ teaspoon 125mg	2 teaspoon 200 mg
60-71 lbs	1 ½ teaspoon 150mg	3 teaspoons 300 mg
72-95 lbs	2 teaspoons 200mg	4 teaspoons 400 mg
96-154lbs	2 teaspoons 200mg	4 teaspoons 400 mg